



10/737,309 (RCE)

FULL ESTIMATED COST	28.79	195.94
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-4.50	-4.50

STN INTERNATIONAL LOGOFF AT 10:24:03 ON 29 AUG 2006

\*\*\*\*\* Welcome to STN International \*\*\*\*\*  
 =>Uploading C:\Program Files\Stnexp\Queries\10737309.str



chain nodes :  
 7 8 9 10 11 12 13 16  
 ring nodes :  
 1 2 3 4 5 6  
 chain bonds :  
 2-8 5-7 7-9 9-11 9-16 10-11 11-13 12-13  
 ring bonds :  
 1-2 1-6 2-3 3-4 4-5 5-6  
 exact/norm bonds :  
 2-8 5-7 7-9 9-11 9-16 10-11 11-13 12-13  
 normalized bonds :  
 1-2 1-6 2-3 3-4 4-5 5-6  
 isolated ring systems :  
 containing 1 :

G1:H,Ak

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:CLASS 9:CLASS 10:CLASS  
 11:CLASS 12:Atom 13:CLASS 16:CLASS

L1 STRUCTURE UPLOADED

=> dis l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1 sam  
L2 3 SEA SSS SAM L1

=> s l1 full  
L3 38 SEA SSS FUL L1


=> file caplus  
=> s l3  
L4 6 L3

=> s l4 and pd<dec 2002  
22748392 PD<DEC 2002  
(PD<20021200)  
L5 4 L4 AND PD<DEC 2002

=> dis l5 1-4 bib abs hitstr

L5 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2003:473270 CAPLUS Full-text  
DN 139:36444  
TI Preparation of substituted ureas as neuropeptide Y5 receptor antagonists  
IN Greenlee, William J.; Huang, Ying; Kelly, Joseph M.; McCombie, Stuart W.;  
Stamford, Andrew W.; Wu, Yusheng  
PA Schering Corporation, USA  
SO U.S. Pat. Appl. Publ., 108 pp., Cont.-in-part of U.S. Ser. No. 950,908.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 2003114517	A1	20030619	US 2002-96390	20020312
	US 6894063	B2	20050517		
	US 2002165223	A1	20021107	US 2001-950908	20010912 <--
	US 2005038100	A1	20050217	US 2004-933016	20040901
PRAI	US 2000-232255P	P	20000914		
	US 2001-950908	A2	20010912		
	US 2002-96390	A3	20020312		
OS	MARPAT 139:36444				
GI					



AB The title compds. [I; Y = II, III; R1 = H, alkyl; R2 = H, alkyl, cycloalkyl, etc.; R3 = IV, V, etc.; j = 0-2; k = 1-2; l = 0-2; m = 0-2; p = 1-3; r = 1-3; R4 = H, OH, halo, etc.; R5 = H, halo, OH, etc.; R6 = alkylSO<sub>2</sub>, cycloalkylSO<sub>2</sub>, heteroarylalkyl, etc.; ], useful as neuropeptide Y5 receptor antagonists for treating obesity, hyperphagia, type II diabetes, insulin resistance, and hypertension, were prepared E.g., a multi-step synthesis of VI, was given. For the compds. I, a range of neuropeptide Y5 receptor binding activity from about 0.2 nM to about 500 nM was observed. Methods of preparing pharmaceutical formulations comprising one or more such compds. I were claimed.


IT 405056-07-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of substituted ureas as neuropeptide Y5 receptor antagonists)

RN 405056-07-7 CAPLUS

CN Urea, N'-(3',5'-difluoro[1,1'-biphenyl]-4-yl)-N-[1-(5-hydroxy-2-pyridinyl)-4-piperidinyl]-N-methyl- (9CI) (CA INDEX NAME)



RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:220568 CAPLUS Full-text

DN 136:263169

TI Preparation of Substituted ureas as neuropeptide Y5 receptor antagonists

IN Greenlee, William J.; Huang, Ying; Kelly, Joseph M.; McCombie, Stuart W.; Stamford, Andrew W.; Wu, Yusheng

PA Schering Corporation, USA

SO PCT Int. Appl., 101 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022592	A2	20020321	WO 2001-US28324	20010912 <--
	WO 2002022592	A3	20020627		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2422013	AA	20020321	CA 2001-2422013	20010912 <--
	AU 2001094547	A5	20020326	AU 2001-94547	20010912 <--
	EP 1322628	A2	20030702	EP 2001-975194	20010912
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	JP 2004509108	T2	20040325	JP 2002-526845	20010912
PRAI	US 2000-232255P	P	20000914		
	WO 2001-US28324	W	20010912		
OS	MARPAT 136:263169				
GI					

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Title compds. [I; A = Q, Q1; R1 = H, F, Cl, CF3, OH; R2 = H, F, Cl, CF3, CN, OCH3, OH; R3 = H, F, Cl, CF3, OCF3, CN, OCH2C6H5, OH; R4 = H, F, Cl; X = NH, NCH3; n = 0, 1, 2; Y = NR5, C:NOH; R5 = SO2CH3, SO2(CH2)2CH3, cyclopropylmethyl, 3-pyridyl, 2-pyridyl, 2-thiazolyl, 2-pyrimidyl, 1-oxo-3-pyridyl, SO2NH2, CH2CONH2, CONH2, NHSO2CH3, SO2(CH2)2OH, C(:NCN)NHCH3, C(:NCN)SCH3, 3-pyridylcarbonyl, cyclobutylcarbonyl, cyclopentylcarbonyl, CON(CH3)2, cyclohexyl; R6 = H, F, Br, Cl, OCH3, OH; R7 = H, F, Cl, OCH3; etc.], stereoisomers, N-oxides, pharmaceutically acceptable salts or hydrates, and prodrugs are disclosed as neuropeptide Y5 receptor antagonists. Method of treating obesity, hyperphagia, type II diabetes, insulin resistance, and hypertension involving title compds. I are claimed. Thus, the title compound II was prepared from N-tert-butoxycarbonyl-4-piperidone, 4-bromophenyl isocyanate, 2-fluorophenylboronic acid, and methanesulfonyl chloride in multiple steps.


IT 405056-07-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of substituted ureas as neuropeptide Y5 receptor antagonists)

RN 405056-07-7 CAPLUS

CN Urea, N'-(3',5'-difluoro[1,1'-biphenyl]-4-yl)-N-[1-(5-hydroxy-2-pyridinyl)-4-piperidinyl]-N-methyl- (9CI) (CA INDEX NAME)



LS ANSWER 3 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1989:90608 CAPLUS Full-text

DN 110:90608

TI Fruit thinning agents containing pyrazoles

IN Kato, Shozo; Noma, Yutaka; Igami, Satoyoshi

PA Tokuyama Soda Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 27 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 63174905	A2	19880719	JP 1987-4945	19870114 <--
	JP 07106964	B4	19951115		
PRAI	JP 1987-4945		19870114		

OS MARPAT 110:90608

GI For diagram(s), see printed CA Issue.

AB Fruit thinning agents contg. title compds. I [R = H, alkyl, (substituted) Ph; R1-R5 = H, halo, (substituted) alkyl, alkoxy, alkylthio, alkoxyalkyl, OH, NO<sub>2</sub>, cyano; R1R2 forms ring; R6 = H, (substituted) alkyl, (substituted) Ph, (substituted) pyridyl; A = CH<sub>2</sub>N; n ≥ 0] as active ingredients are described. A solution of 5-amino-1,3-dimethylpyrazole in C<sub>6</sub>H<sub>6</sub> was treated with 2,4-MeClC<sub>6</sub>H<sub>3</sub>OCHMeCOCl to give 84.8% N-pyrazolylpropanamide derivative II, which at 200 ppm showed fruit thinning to 23.8% in mandarin orange. A wettable powder was formulated containing II 10, polyoxyethylene nonylphenyl ether 2, clay 40, and zeolite 48 weight parts.


IT 118912-52-0P


RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of, as fruit thinning agent)

RN 118912-52-0 CAPLUS

CN Propanamide, 3-(2,4-dichlorophenoxy)-N-[1-(5-hydroxy-2-pyridinyl)-3,5-dimethyl-1H-pyrazol-4-yl]- (9CI) (CA INDEX NAME)



PAGE 1-A





L5 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2006 ACS on STN  
 AN 1987:576028 CAPLUS Full-text  
 DN 107:176028  
 TI Preparation of [(phenoxyalkanoyl)amino]pyrazole derivatives as herbicides, fungicides and bactericides  
 IN Kato, Shozo; Takematsu, Tetsuo; Igami, Satoyoshi; Ogasawara, Masaru  
 PA Tokuyama Soda Co., Ltd., Japan  
 SO Jpn. Kokai Tokkyo Koho, 18 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 62138475	A2	19870622	JP 1985-277887	19851212 <--
	JP 05080469	B4	19931109		
PRAI	JP 1985-277887		19851212		
GI					




AB The title compds. I [R1-R5 = H, halo, (un)substituted alkyl, alkoxy(alkyl), alkylthio, OH, NO<sub>2</sub>, cyano, or R1R2 being adjacent and completing a fused ring; R6 = (un)substituted alkyl, Ph or pyridyl; n = integer], useful as herbicides, fungicides and bactericides, were prepared A solution of 0.0042 mmol 2,4-Cl<sub>2</sub>C<sub>6</sub>H<sub>3</sub>OCH<sub>2</sub>COC<sub>2</sub>H<sub>5</sub> in benzene was added dropwise to a solution of 0.0032 mmol 5-amino-1,3-dimethylpyrazole and 0.0042 mmol Et<sub>3</sub>N in benzene and the mixture was stirred overnight to give 0.85 g a pyrazole derivative II (R7 = H, R8 = Me). In preemergence period, I at 100g/10 are controlled by 90-100% various weeds, e.g., Scirpus juncooides. II (R7 = Cl, R8 = 3,4-dichlorophenyl) in vitro is active against fungi, e.g., Pellicularia sasakii and a bacterium, Staphylococcus aureus.

IT 110731-75-4P  
 RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of, as herbicide, fungicide and bactericide)  
 RN 110731-75-4 CAPLUS  
 CN Acetamide, 2-(2,4-dichlorophenoxy)-N-[1-(5-hydroxy-2-pyridinyl)-3,5-dimethyl-1H-pyrazol-4-yl]- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



=&gt; s l4 not l5

L6 2 L4 NOT L5

=&gt; dis l6 1-2 bib abs

L6 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:238237 CAPLUS Full-text

DN 144:311912

TI Preparation of arylpyridines as inhibitors of hedgehog signalling.

IN Gunzner, Janet; Sutherlin, Daniel; Stanley, Mark; Bao, Liang; Castanedo, Georgette; Lalonde, Rebecca; Wang, Shumei; Reynolds, Mark; Savage, Scott; Malesky, Kimberly; Dina, Michael

PA Genentech, Inc., USA; Curis Incorporation


SO PCT Int. Appl., 256 pp.



CODEN: PIXXD2

DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2006028958	A2	20060316	WO 2005-US31284	20050902
	WO 2006028958	A3	20060413		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM US 2006063779 A1 20060323 US 2005-217663 20050902 PRAI US 2004-607367P P 20040902 OS MARPAT 144:311912 GI				



AB Title compds. [I; A = carbocyclyl, heterocyclyl; X = alkylene, NR<sub>4</sub>CO, NR<sub>4</sub>CS, NR<sub>4</sub>SO<sub>2</sub>, NR<sub>4</sub>PO(OH), etc.; R<sub>1</sub> = (substituted) alkyl, carbocyclyl, heterocyclyl; R<sub>2</sub> = halo, OH, (substituted) alkyl, acyl, alkoxy; R<sub>3</sub> = halo, OH, CO<sub>2</sub>H, (substituted) alkyl, acyl, alkoxy, alkoxy carbonyl, carbamoyl, alkylthio, sulfinyl, sulfonyl, carbocyclyl, heterocyclyl; R<sub>4</sub> = H, alkyl; m, n = 0-3], were prepared for treatment of cancer (no data). Thus, N-[4-chloro-3-(pyridin-2-yl)phenyl]-6-chloro-3-carboxamide and 2-morpholinoethylamine were heated in BuOH in a sealed tube to give 6-(2-morpholinoethylamino)-N-[4-chloro-3-(pyridin-2-yl)phenyl]-6-chloro-3-carboxamide.

L6 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:531358 CAPLUS Full-text

DN 141:89014

TI Preparation of pyridylcyclohexyl phenylpropanamide derivatives as NR2B receptor antagonists

IN Kawai, Makoto; Nakamura, Hiroshi; Shimokawa, Hirohisa

PA Pfizer Japan Inc., Japan; Pfizer Inc.

SO PCT Int. Appl., 91 pp.

CODEN: PIXXD2


DT Patent  
LA English  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004054579	A1	20040701	WO 2003-IB5757	20031205
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				


## 10/737,309 (RCE)

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,  
 GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
 LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ,  
 OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,  
 BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
 ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
 TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

CA 2510322 AA 20040701 CA 2003-2510322 20031205  
 AU 2003303042 A1 20040709 AU 2003-303042 20031205  
 EP 1575586 A1 20050921 EP 2003-813240 20031205  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 BR 2003017409 A 20051108 BR 2003-17409 20031205  
 JP 2006511528 T2 20060406 JP 2004-560052 20031205  
 US 2004152715 A1 20040805 US 2003-737309 20031216  
 PRAI US 2002-434361P P 20021217  
 WO 2003-IB5757 W 20031205  
 OS MARPAT 141:89014  
 GI



I



II

AB Title compds. I [wherein R2 = H or OH; or R forms a covalent bond with ring A; R3 = H or alkyl; R4 = (un)substituted (hetero)aryl; R5 = OH or alkylsulfonylamino; R6 = H, halo, alkylalkoxy; A = cycloalkylene; X = a covalent bond, alkylene, (hetero)alkenylene, etc.; Z = C or N; and pharmaceutically acceptable ester or salts thereof] were prepared as. For example, II•HCl was given in 5-step synthesis starting from trans-4-aminocyclohexanol and 3-phenylpropanoic acid. I showed Ki values from 2.7  $\mu$ M to 8.9  $\mu$ M with respect to inhibition of binding at the NR2B receptor. Thus, I and their pharmaceutical compns. are useful for the treatment of disease conditions caused by over activation of NMDA NR2B receptor such as pain, or the like in mammals.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> log

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y

COST IN U.S. DOLLARS

SINCE FILE  
ENTRY

TOTAL  
SESSION